

UCLA

UCLA Previously Published Works

Title

Using hybrid transcription factors to study gene function in rice.

Permalink

<https://escholarship.org/uc/item/94r8x2ct>

Journal

Science China. Life sciences, 58(11)

ISSN

1674-7305

Authors

Zhao, Tao
Liu, Jun
Li, Hong-Yu
[et al.](#)

Publication Date

2015-11-01

DOI

10.1007/s11427-015-4937-x

Peer reviewed

Using hybrid transcription factors to study gene function in rice

ZHAO Tao^{1†}, LIU Jun^{1†}, LI Hong-Yu^{1†}, LIN Jian-Zhong^{2†}, BIAN Ming-Di^{3†}, ZHANG Chun-Yu¹, ZHANG Yong-Xing¹, PENG Yu-Chong², LIU Bin^{1*} & LIN ChenTao⁴

¹Institute of Crop Sciences, Chinese Academy of Agricultural Sciences, Beijing 100081, China;

²College of Biology, Hunan Province Key Laboratory of Plant Functional Genomics and Developmental Regulation, Hunan University, Changsha 410082, China;

³Laboratory of Soil and Plant Molecular Genetics, College of Plant Science, Jilin University, Changchun 130062, China;

⁴Department of Molecular, Cell & Developmental Biology, University of California, Los Angeles, CA90095, USA

Received September 14, 2015; accepted September 18, 2015; published online September 24, 2015

Citation: Zhao T, Liu J, Li HY, Lin JZ, Bian MD, Zhang CY, Zhang YX, Peng YC, Liu B, Lin CT. Using hybrid transcription factors to study gene function in rice. *Sci China Life Sci*, 2015, 58: 1160–1162, doi: 10.1007/s11427-015-4937-x

Dear Editors,


Transcription factors (TF) play important roles in plant growth and development, which determine crop productivity, quality, and stress tolerance. However, the basic mechanism governing transcription regulation by TF remains largely unclear in plants. It is well known that TF proteins contain at least two functional domains, the DNA-binding domain and the transcription regulatory domain. The VP16 transcription activation domain of human Herpes simplex virus and the EAR transcription suppression domain of many plant TF proteins are two of the best-known transcription regulatory domains [1,2]. It has been reported that both VP16 and EAR domains exert their respective transcription activation/repression functions in heterologous protein and/or cellular environment [2–4], supporting a hypothesis that the transcription regulatory domains function independent of other functional domains of TF proteins. However, this hypothesis has not been examined systematically. We reasoned that a phenotypic analysis of transgenic expression of all or most TF genes from a plant species in the form of fusion proteins, which are composed of the coding sequences of TFs fused to the VP16 or EAR domain, would allow a systematic test of this hypothesis, and we reported

here the overall result of such an experiment.

We cloned cDNAs for 1,685 out of the 3,148 predicted transcription factor genes from rice (*Oryza sativa subsp. Japonica*) (Figure 1A, Table S1 in Supporting Information), and prepared 2,980 HTF Ti plasmid constructs that express the EAR4 (tetrameric repeats of EAR)-TF fusion proteins or the VP64 (tetrameric repeats of VP16)-TF fusion proteins under the control of the maize ubiquitin promoter [5] (Figure 1B, Table S2 in Supporting Information). We obtained 57,751 independent transgenic lines in the *Japonica* rice accession Kitaake that express a total of 2,470 HTFs constructs. On average, 23 independent transgenic lines were generated per construct (Figure 1C, Table S3 in Supporting Information). Approximately 13% (7,719/57,751 events) of transgenic lines showed phenotypic alterations in comparison to the wild-type parents, including changes in morphology, heading date, or grain yield (Figure S1 in Supporting Information). Expression of 957 HTF fusion proteins, including 343 EAR4 HTF and 614 VP64 HTF proteins, caused the phenotypic alteration in at least three independent transgenic lines, and they were investigated in more details (Figure 1D). 132 pairs of these 957 HTFs exhibited phenotypic alterations in both the EAR4 and VP64 configurations, of which 86, 30 and 16 pairs of EAR4 and VP64 HTFs resulted in similar, different, or opposite phenotypes, respectively (Figure 1D, Pie diagram). It is surprising that only 12% (16/132) of EAR4- and VP64-fusion

[†]Contributed equally to this work

*Corresponding author (email: liubin05@caas.cn)



On the other hand, those 16 HTFs that conferred opposite phenotypes in the VP64 or EAR4 configuration provide a relatively simpler explanation for the mode of action of the respective rice TF genes (Table S4 in Supporting Information). For example, a MYB-like transcription factor (encoded by *LOC_Os06g45890*), referred to as TF865 in this study, showed opposite growth phenotype in the EAR4 or VP64 configuration. Transgenic lines expressing the VP64-fusion proteins (*CV865*) exhibited dwarf phenotype, whereas those expressing the EAR4-fusion proteins (*CE865*) grew significantly taller than the wild-type control (Figure 1E and 1H). Overexpression of both *CV865* and *CE865* transgenes were confirmed by the quantitative reverse transcription PCR (qRT-PCR) or the immunoblot

analyses (Figure 1F and 1G). In comparison to that of the wild-type control, the grain yield per plant of transgenic lines expressing the EAR4-fusion protein (CE865) or the VP64-fusion protein (CV865) increased (50%–100%) or decreased (10%–20%), respectively (Figure 1I). These results are consistent with the hypothesis that this MYB-like TF gene may regulate rice growth and development or metabolism to affect grain yield. Further study of other HTF that caused various phenotypic variations will likely reveal the functional roles and modes of action for other rice TF genes. Transcription factors regulate almost all aspects of biological processes by dynamically modulating target gene transcription. A large percentage of genes, ranged from 1%–6% in different Eukaryotic species, encode TFs [10]. Only a relatively small fraction of eukaryotic TFs' DNA sequence preferences are experimentally determined, but their biological functions are much less determined. The functional redundancy of many plant TF genes imposes a major technical difficulty in the study of physiological roles of many TF genes. The HTF transgenic resource reported here provides a valuable tool complement to the loss-of-function resources reported previously for the functional study of rice genes.

This work is supported by the MOA Transgenic Research Grant (2010ZX08010-002 to ICS-CAAS), Agricultural Science and Technology Innovation Program of CAAS, and National Natural Science Foundation of China (31422041, 31371649, 31301298 and 31170172).

- 1 Sadowski I, Ma J, Triezenberg S, Ptashne M. GAL4-VP16 is an unusually potent transcriptional activator. *Nature*, 1988, 335: 563–564
- 2 Kagale S, Rozwadowski K. EAR motif-mediated transcriptional repression in plants: an underlying mechanism for epigenetic regulation of gene expression. *Epigenetics*, 2011, 6: 141–146
- 3 Ohta M, Matsui K, Hiratsu K, Shinshi H, Ohme-Takagi M. Repression domains of class II ERF transcriptional repressors share an essential motif for active repression. *Plant Cell*, 2001, 13: 1959–1968
- 4 Herve C, Dabos P, Bardet C, Jauneau A, Auriac MC, Ramboer A, Lacout F, Tremousaygue D. In vivo interference with AtTCP20 function induces severe plant growth alterations and deregulates the expression of many genes important for development. *Plant Physiol*, 2009, 149: 1462–1477
- 5 Cornejo MJ, Luth D, Blankenship KM, Anderson OD, Blechl AE. Activity of a maize ubiquitin promoter in transgenic rice. *Plant Mol Biol*, 1993, 23: 567–581
- 6 Wysocka J, Herr W. The herpes simplex virus VP16-induced complex: the makings of a regulatory switch. *Trends Biochem Sci*, 2003, 28: 294–304
- 7 Xiao H, Pearson A, Coulombe B, Truant R, Zhang S, Regier JL, Triezenberg SJ, Reinberg D, Flores O, Ingles CJ, et al. Binding of basal transcription factor TFIID to the acidic activation domains of VP16 and p53. *Mol Cell Biol*, 1994, 14: 7013–7024
- 8 Hall DB, Struhl K. The VP16 activation domain interacts with multiple transcriptional components as determined by protein-protein cross-linking in vivo. *J Biol Chem*, 2002, 277: 46043–46050
- 9 Wang L, Kim J, Somers DE. Transcriptional corepressor TOPLESS complexes with pseudoresponse regulator proteins and histone deacetylases to regulate circadian transcription. *Proc Natl Acad Sci USA*, 2013, 110: 761–766
- 10 Jin J, Zhang H, Kong L, Gao G, Luo J. PlantTFDB 3.0: a portal for the functional and evolutionary study of plant transcription factors. *Nucleic Acids Res*, 2014, 42: D1182–1187

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Supporting Information

Figure S1 The number of HTFs in EAR4 or VP64 configuration showing indicated phenotypic alterations.

Table S1 The list of 1685 transcription factor genes cloned into pDONR vector.

Table S2 The list of 2980 HTF constructs.

Table S3 The numbers of independent transgenic events of each HTF construct.

Table S4 The list of 16 pairs of EAR4- and VP64-fusion proteins causing opposite phenotypes.

The supporting information is available online at life.scichina.com and link.springer.com. The supporting materials are published as submitted, without typesetting or editing. The responsibility for scientific accuracy and content remains entirely with the authors.